

Matrix Model for Discretized Moduli Space

L. Chekhov*

*Steklov Mathematical Institute
Vavilov st.42, GSP-1, 117966 Moscow, Russia*

Abstract

We study the algebraic geometrical background of the Penner–Kontsevich matrix model with the potential $N\alpha \operatorname{tr}(-\frac{1}{2}\Lambda X \Lambda X + \log(1 - X) + X)$. We show that this model describes intersection indices of linear bundles on the discretized moduli space right in the same fashion as the Kontsevich model is related to intersection indices (cohomological classes) on the Riemann surfaces of arbitrary genera. The special role of the logarithmic potential originated from the Penner matrix model is demonstrated. The boundary effects which was unessential in the case of the Kontsevich model are now relevant, and intersection indices on the discretized moduli space of genus g are expressed through Kontsevich's indices of the genus g and of the lower genera.

*E-mail: chekhov@qft.mian.su

1 Introduction.

The last three years of development in matrix models initiated by the papers [1] revealed a lot of applications of these models in various branches of mathematical physics: two-dimensional quantum field theory, intersection theory on the moduli space of Riemann surfaces, integrable hierarchies, matrix integrals, random surfaces and others. The approach by [1] where the explicit solution was presented deal with triangulated Riemann surfaces where any triangulation determines some singular metric obtained by the arrangement of equilateral triangles. One can think that when the number of triangles tends to infinity these singular metrics approximate “random metrics” on the surface. These triangulations were presented by a hermitean $N \times N$ one-matrix model

$$\int \exp(\operatorname{tr} P(X)) DX, \quad (1.1)$$

where $P(X) = \sum_n T_n \operatorname{tr} X^n$, T_n being times for the one-matrix model. For this system discrete Toda chain equations holds with an additional Virasoro symmetry imposed [2]. In the limit $N \rightarrow \infty$ the Kortevég-de-Vries equation arises. The partition function of the two-dimensional gravity for this approach is a series in an infinite number of variables and coincides with the logarithm of some τ -function for KdV hierarchy.

Another approach to the two-dimensional gravity is to do the integral over all classes of conformally equivalent metrics on Riemann surfaces. It may be presented as an integral over the finite-dimensional space of conformal structures. This integral has a cohomological description as an intersection theory on the compactified moduli space of complex curves. Edward Witten presented compelling evidence for a relationship between random surfaces and the algebraic topology of moduli space [3], [4]. In fact, he suggested that these expressions coincide since both satisfy the same equations of KdV hierarchy. It was Maxim Kontsevich who proved this assumption [5]. Surprisingly, he explicitly presented a new matrix model defining exactly the values of intersection indices or, on the language of 2D gravity, correlation functions of observables $\mathcal{O}_n$ of the type

$$\langle \mathcal{O}_{n_1} \dots \mathcal{O}_{n_s} \rangle_g, \quad (1.2)$$

where $\langle \dots \rangle_g$ denotes the expectation value on a Riemann surface with g handles. Then the string partition function $\tau(t)$ has an asymptotic expansion of the form

$$\tau(t) = \exp \sum_{g=0}^{\infty} \left\langle \exp \sum_n t_n \mathcal{O}_n \right\rangle_g, \quad (1.3)$$

and it is a tau-function of the KdV hierarchy taken at a point of Grassmannian where it is invariant under the action of the set of the Virasoro constraints: $\mathcal{L}_n \tau(t) = 0$, $n \geq -1$ [6], [7], [8], [9]. One might say that the Kontsevich model is used to triangulate moduli space, whereas the original models triangulated Riemann surfaces (see e.g. [10]).

The generalization of the Kontsevich model — so-called Generalized Kontsevich Model (GKM) [11] is related to the two-dimensional Toda lattice hierarchy and it originated from

the external field problem defined by the integral

$$\mathcal{Z}[\Lambda; N] = \int DX \exp \{ N \operatorname{tr} (\Lambda X - V_0(X)) \}, \quad (1.4)$$

where $V_0(X) = \sum_n t_n \operatorname{tr} X^n$ is some potential, t_n are related to times of the hierarchy. This model is equivalent to the Kontsevich integral for $V_0(X) \sim \operatorname{tr} X^3$. To solve the integral (1.4) one may use the Schwinger–Dyson equation technique [12] written in terms of eigenvalues of Λ . The Kontsevich model was solved in the genus expansion in the papers [8], [13] for genus zero (planar diagrams) and in [14] for higher genera.

Recently, the Kontsevich–Penner model was introduced [15]. The Lagrangian of this model has the following form:

$$\mathcal{Z}[\Lambda] = \int DX \exp \left(N \operatorname{tr} \left\{ -\frac{1}{2} \Lambda X \Lambda X + \alpha [\log(1 + X) - X] \right\} \right), \quad \Lambda = \operatorname{diag}(\Lambda_1, \dots, \Lambda_N). \quad (1.5)$$

This model may be readily reduced to (1.4) with $V_0(X) = -X^2/2 + \alpha \log X$. It was solved in genus expansion in [15], [16]. It was shown in [17], [18] that it is in fact equivalent to the one-matrix hermitean model (1.1) with the general potential

$$P(X) = \sum_{n=0}^{\infty} T_n \operatorname{tr} X^n, \quad (1.6)$$

which times are defined by the kind of Miwa transform:

$$T_n = \frac{1}{n} \operatorname{tr} \Lambda^{-n} - \frac{N}{2} \delta_{n2} \quad \text{for } n \geq 1 \quad \text{and} \quad T_0 = \operatorname{tr} \log \Lambda^{-1}. \quad (1.7)$$

Thus the Kontsevich–Penner model may be treated as an intermediate link between 2D gravity described by the Kontsevich model and random surface technique.

Indeed, as we shall demonstrate, this new model describes in a very natural way the Kontsevich indices for the case of discretized moduli space. If we do not cast the moduli space boundary effects (which are relevant in this model), then the coefficients of expansion are just the Kontsevich indices, but since we deal with the closure of the moduli space, the answer is tuned in a way to incorporate the Riemann surfaces which are boundary components under the Deligne and Mumford reduction procedure [19]. Then when taking the scaling limit (but keeping N finite), we get just the Kontsevich model. From the other hand, taking another rescaling we obtain the Penner model describing virtual Euler characteristics of the moduli space via the cell decomposition. Thus, this model provides a bridge between Harer, Zagier and Penner theory [20], [21] describing virtual Euler characteristics on moduli space and the Kontsevich theory giving intersection numbers of stable cohomology classes on the moduli space.

2 The geometric approach to the Kontsevich model.

In his original paper [5] Kontsevich proved that

$$\sum_{d_1, \dots, d_s=0}^{\infty} \langle \tau_{d_1}, \tau_{d_2}, \dots, \tau_{d_s} \rangle \prod_{i=1}^s (2d_i - 1)!! \lambda_i^{-(2d_i+1)} = \sum_{\Gamma} \frac{2^{-n_0}}{\# \operatorname{Aut}(\Gamma)} \prod_{\{ij\}} \frac{2}{\lambda_i + \lambda_j}, \quad (2.1)$$

where the objects standing in angular brackets on the left-hand side are (rational) numbers describing intersection indices, and on the right-hand side the sum runs over all oriented connected trivalent “fatgraphs” Γ with s labeled boundary components, regardless of the genus, n_0 is the number of vertices of Γ , the product runs over all the edges in the graph and $\# \text{Aut}$ is the volume of discrete symmetry group of the graph Γ .

The amazing result by Kontsevich is that the quantity on the right hand side of (2.1) is equal to a free energy in the following matrix model:

$$e^{F_N(\Lambda)} = \frac{\int dX \exp\left(-\frac{1}{2} \text{tr} \Lambda X^2 + \frac{i}{6} \text{tr} X^3\right)}{\int dX \exp\left(-\frac{1}{2} \text{tr} \Lambda X^2\right)}, \quad (2.2)$$

where X is an $N \times N$ hermitian matrix and $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_N)$. The distinct feature of the expression (2.1) is that in spite of the fact that each selected diagram has quantities $(\lambda_i + \lambda_j)$ in the denominator, when taking a sum over all diagrams of the same genus and the same number of boundary components all these quantities are cancelled with the ones from nominator.

Feynman rules for the Kontsevich matrix model are the following: as in the usual matrix models, we deal with so-called “fat graphs” or “ribbon graphs” with propagators having two sides, each carries corresponding index. The Kontsevich model varies from the standard one-matrix hermitian model since there appear additional variables λ_i associated with index loops in the diagram, the propagator being equal to $2/(\lambda_i + \lambda_j)$, where λ_i and λ_j are variables of two cycles (perhaps the same cycle) which the two sides of propagator belong to. Also there are trivalent vertices presenting the cell decomposition of the moduli space.

It is instructive to consider the simplest example of genus zero and three boundary components which we symbolically label λ_1 , λ_2 and λ_3 . There are two kinds of diagrams giving the contribution in this order (Fig.1). The contribution to the free energy arising from this sum is

$$\begin{aligned} & \frac{1}{6(\lambda_1 + \lambda_2)(\lambda_1 + \lambda_3)(\lambda_2 + \lambda_3)} + \frac{1}{3} \left\{ \frac{1}{4\lambda_1(\lambda_2 + \lambda_1)(\lambda_3 + \lambda_1)} + \right. \\ & + \left. (1 \rightarrow 2, 2 \rightarrow 3, 3 \rightarrow 1) + (1 \rightarrow 3, 3 \rightarrow 2, 2 \rightarrow 1) \right\} \\ & = \frac{2\lambda_1\lambda_2\lambda_3 + \lambda_2\lambda_3(\lambda_2 + \lambda_3) + \lambda_1\lambda_3(\lambda_1 + \lambda_3) + \lambda_1\lambda_2(\lambda_1 + \lambda_2)}{12\lambda_1\lambda_2\lambda_3(\lambda_1 + \lambda_2)(\lambda_1 + \lambda_3)(\lambda_2 + \lambda_3)} \\ & = \frac{1}{12\lambda_1\lambda_2\lambda_3}. \end{aligned} \quad (2.3)$$

This example demonstrates the cancellations of $(\lambda_i + \lambda_j)$ -terms in the denominator above mentioned.

Figure 1. the $g=0, s=3$ contribution to Kontsevich's model

Now the sketch of Kontsevich's proof is in order. Let us associate with each edge e_i of a fat graph its length $l_i > 0$. We consider the orbispace $\mathcal{M}_{g,n}^{comb}$ of fat graphs with all possible lengths of edges and arbitrary valencies of vertices. Two graphs are equivalent if an isomorphism between them exists. Let us introduce an important object — the space of $(2,0)$ -meromorphic differentials $\omega(z)dz^2$ on a Riemann surface with g handles and n punctures, the only poles of $\omega(z)$ are n double poles placed in the points of punctures with strictly positive quadratic residues $p_i^2 > 0$, ($i = 1, \dots, n$). It is Strebel's theorem [22] which claims that the natural mapping from $\mathcal{M}_{g,n}^{comb}$ to the moduli space $\mathcal{M}_{g,n} \times \mathbf{R}_+^n$, where $\mathbf{R}_+^n$ is the space of residues, $p_i > 0$ being perimeters of cycles, is homeomorphism. Thus, varying l_j and taking the composition of all graphs we span the whole space $\mathcal{M}_{g,n} \times \mathbf{R}_+^n$.

Each cycle can be interpreted as a boundary components I_i of the Riemann surface since in the Strebel metric it can be presented as half-infinite cylinder with the puncture point placed at infinity. The boundary of it consists of a finite number of intervals (edges). We consider a set of line bundles $\mathcal{L}_i$ which fiber at a point $\Sigma \in \mathcal{M}_{g,n}$ is the cotangent space to the puncture point x_i on the surface Σ . The first Chern class of the line bundle $\mathcal{L}_i$ admits a representation in terms of the lengths of the intervals l_j . The perimeter of the boundary component is $p_i = \sum_{l_\alpha \in I_i} l_\alpha$ and

$$c_1(\mathcal{L}_i) = \sum_{\substack{a,b \in I_i \\ a < b}} d \left(\frac{l_a}{p_i} \right) \wedge d \left(\frac{l_b}{p_i} \right), \quad (2.4)$$

where the cyclic ordering is assumed. Following Kontsevich we introduce 2-form Ω :

$$\Omega = \sum_{i=1}^n \frac{1}{2} p_i^2 c_1(\mathcal{L}_i). \quad (2.5)$$

The intersection indices are generated by the integrals over appropriate power $d = 3g - 3 + n$ of the form Ω :

$$\begin{aligned} \frac{2^d}{d!} \int_{\mathcal{M}_{g,n}} \Omega^d &= \frac{1}{d!} \int_{\mathcal{M}_{g,n}} (p_1^2 c_1(\mathcal{L}_1) + \dots + p_n^2 c_1(\mathcal{L}_n))^d = \\ &= \sum_{\sum d_i = d} \prod_{i=1}^n \frac{p_i^{2d_i}}{d_i!} \langle \tau_{d_1} \dots \tau_{d_n} \rangle_g. \end{aligned} \quad (2.6)$$

One important note is in order. It is a theorem by Kontsevich that these integrations extend continuously to the closure of the moduli space $\bar{\mathcal{M}}_{g,n}$ following the procedure by Deligne and Mumford [19], and the proper integration goes over $\bar{\mathcal{M}}_{g,n} \times \mathbf{R}_+^n$. (It means that we deal with a stable cohomological class of curves.)

Taking the Laplace transform over variables p_i we get:

$$\int_0^\infty dp_i e^{-p_i \lambda_i} p_i^{2d_i} = (2d_i)! \lambda_i^{-2d_i-1} \quad (2.7)$$

for the quantities standing on the right-hand side of (2.6). On the left-hand side we have

$$\int_0^\infty \dots \int_0^\infty dp_1 \wedge \dots \wedge dp_n e^{-\sum p_i \lambda_i} \int_{\mathcal{M}_{g,n}} e^\Omega, \quad (2.8)$$

and due to cancellations of all p_i^2 multipliers with p_i 's in denominators of the form Ω we get:

$$e^\Omega dp_1 \wedge \dots \wedge dp_n = c \prod_\alpha \wedge dl_\alpha. \quad (2.9)$$

Surprisingly, the constant c does depend only on Euler characteristic of the graph Γ , $c = 2^{-\kappa}$, $\kappa = 2 - 2g = \# \text{ vertices} - \# \text{ edges} + n$. Thus we have

$$\sum_{\sum d_i = d} \prod_{i=1}^n \frac{(2d_i - 1)!!}{\lambda_i^{2d_i+1}} < \tau_{d_1} \dots \tau_{d_n} >_g = \sum_\Gamma \frac{2^{-\kappa}}{\# \text{Aut } \Gamma} \int [dl] \exp \left(\sum_\alpha l_\alpha (\lambda_\alpha^{(1)} + \lambda_\alpha^{(2)}) \right). \quad (2.10)$$

Here $\lambda_\alpha^{(1)}$ and $\lambda_\alpha^{(2)}$ are variables of two cycles divided by α th edge. Integration over all dl_α gives us eventually the relation (2.1).

3 The Penner–Kontsevich model.

Now let us turn to the case of the Penner–Kontsevich model (PK model). It includes in variance with the Kontsevich model all powers of X^n in the potential since it describes the partition of moduli space into cells of a simplicial complex, the sum running over all simplices with different dimensions. (On the language of the Kontsevich model the lower the dimension, the more and more edges of the fat graph are reduced). Then the virtual Euler characteristic is obtained by weighting the simplices by

$$\frac{(-1)^{d_F}}{|G_F|} \quad (3.1)$$

where $|G_F|$ denotes the order of a stabilizer of the subgroup of the mapping class group, that is, the order of the symmetry group of the corresponding fat graph. It is the Penner model which gives the answer for the sum over F in (3.1) as a free energy for a matrix model [21],[24]:

$$\sum_F N^{2-2g} t^{2-2g-n} \frac{(-1)^{d_F}}{|G_F|} = \log \int dX e^{Nt \text{tr} [\log(1-X) + X]}, \quad (3.2)$$

where n is a number of punctures on genus g Riemann surface. Expansion of the free energy of this model in N and t reveals logarithmic corrections which is a feature of $c = 1$ theories.

We find the Feynman rules for the Kontsevich–Penner theory (1.5). First, as in the standard Penner model, we have vertices of all orders in X . Due to rotational symmetry, the factor $1/n$ standing with each X^n cancels, and only symmetrical factor (3.1) survives. Also there is a factor $(-\alpha)$ standing with each vertex. As in Kontsevich model, there are

variables λ_i associated with each cycle. But the form of propagator changes — instead of $2/(\lambda_i + \lambda_j)$ we have $1/(\lambda_i \lambda_j + \alpha)$.

Let us consider the same case ($g = 0, n = 3$) as for Kontsevich model. One additional diagram resulting from vertex X^4 arises and gives the contribution with opposite sign (Fig.2).

Fig.2. $g=0, s=3$ contribution to Penner–Kontsevich’s model.

This contribution is (symmetrized over λ_1, λ_2 and λ_3):

$$\begin{aligned}
& - \frac{1}{3} \left\{ \frac{\alpha}{2(\lambda_1 \lambda_2 + \alpha)(\lambda_1 \lambda_3 + \alpha)} - \text{perm.} \right\} + \frac{\alpha^2}{6(\lambda_1 \lambda_2 + \alpha)(\lambda_1 \lambda_3 + \alpha)(\lambda_2 \lambda_3 + \alpha)} \\
& + \frac{1}{3} \left\{ \frac{\alpha^2}{2(\lambda_1^2 + \alpha)(\lambda_1 \lambda_2 + \alpha)(\lambda_1 \lambda_3 + \alpha)} + \text{perm.} \right\} \quad (3.3)
\end{aligned}$$

Again collecting all terms we get:

$$\frac{\alpha}{6 \prod_{i < j} (\lambda_i \lambda_j + \alpha)} \left\{ - \sum_{i < j} \lambda_i \lambda_j - 2\alpha + \alpha \left(\frac{\lambda_2 \lambda_3 + \alpha}{\lambda_1^2 + \alpha} + \frac{\lambda_1 \lambda_2 + \alpha}{\lambda_3^2 + \alpha} + \frac{\lambda_1 \lambda_3 + \alpha}{\lambda_2^2 + \alpha} \right) \right\}, \quad (3.4)$$

and after a little algebra we obtain an answer:

$$F_{0,3} = \alpha \frac{-\lambda_1 \lambda_2 - \lambda_1 \lambda_3 - \lambda_2 \lambda_3 + \alpha}{3(\lambda_1^2 + \alpha)(\lambda_2^2 + \alpha)(\lambda_3^2 + \alpha)}. \quad (3.5)$$

We see that here, just as in standard Kontsevich model, the cancellation of intertwining terms in the denominator occurs that leads to factorization of the answer over $1/(\lambda_i^2 + \alpha)$ -terms. It should reveal an underlying geometric structure of the model under consideration, where quantities like (2.1) are expected to arise.

4 Relation between KP model and discretized moduli space.

Now we turn to the description of underlying differential–geometric structure of the Kontsevich–Penner model. Let us consider the case of a discretized moduli space. Its description can be done most properly in terms of lengths of edges constituting boundary

components of the surface. We assume all these lengths to be integers (probably zeros) scaled by some factor ϵ . Thus, perimeters p_i belong now to $\epsilon \cdot \mathbf{Z}_+$. Then we can present the first Chern class of the line bundle $\mathcal{L}_i$ in the same form as above

$$c_1(\mathcal{L}_i) = \sum_{\substack{a,b \in I_i \\ a < b}} d \left(\frac{n_a}{p_i} \right) \wedge d \left(\frac{n_b}{p_i} \right), \quad (4.1)$$

where dn_a are symbols satisfying standard relations, dn_a lie in the cotangent space to the continuous moduli space taken at a point $(n_1, \dots, n_{6g-6+3n})$ which belongs also to the discretized moduli space. The action of the external derivative d follows the same rules as in continuous case and the integration is replaced by a discrete half-infinite sum. The 2-form Ω is defined by the same formula (2.5), thus the “volume formula” (2.9) is also preserved. Subtleties appear when one has to integrate over discrete moduli space $\mathcal{M}_{g,n}^{disc}$. First, there are points in $\mathcal{M}_{g,n}^{disc}$ which do not lie on the boundary of the moduli space but in which some of the lengths n_a are equal zero. These points correspond to graphs containing vertices of order greater than three. In the continuum limit we did not take into account such graphs since they correspond to subdomains of lower dimensions in the interior of the moduli space and because the integration measure is continuous we may neglect them. Now the situation changed and we should cast these diagrams as well.

Second, now we should explicitly take into account curves which are reduced by Deligne–Mumford procedure [19]. Fortunately, we can present an explicit integration over the boundary $\partial \mathcal{M}_{g,n}^{disc}$ of the discrete moduli space because it expands into a sum over direct products of connected components of lower genera and a number of additionally inserted punctures corresponding to reduced handles of the surface. Doing all possible reductions we span the whole closure of $\mathcal{M}_{g,n}^{disc}$:

$$\frac{2^d}{d!} \int_{\mathcal{M}_{g,n}^{disc}} \Omega^d = \frac{1}{d!} \int_{\bar{\mathcal{M}}_{g,n}^{disc}} \left(\sum_{i=1}^n p_i^2 c_1(\mathcal{L}_i) \right)^d - \frac{1}{d!} \int_{\partial \bar{\mathcal{M}}_{g,n}^{disc}} \left(\sum_{i=1}^n p_i^2 c_1(\mathcal{L}_i) \right)^{d'}. \quad (4.2)$$

Here $d' < d$ depends on the power of reduction.

A point $\Sigma \in \partial \mathcal{M}_{g,n}^{disc}$ is a union of s ($1 \leq s \leq n + 2g - 2$) connected components Σ_{g_j, n_j, k_j} , ($j = 1, \dots, s$). Each surface Σ_{g_j, n_j, k_j} has genus g_j , $\sum_{j=1}^s g_j \leq g$, n_j original punctures ($\sum_{j=1}^s n_j = n$) and k_j additional punctures $\mathcal{P}_l^{(j)}$ arising from the reduction procedure. The linear bundles $\mathcal{L}_i$ are associated with n_j points of this surface but not with the new $\mathcal{P}_l^{(j)}$. Explicitly the boundary $\partial \mathcal{M}_{g,n}^{disc}$ can be presented as a finite set of disconnected pieces, each of them is in its turn the direct product of lower dimensional closed moduli spaces weighted with $(-1)^{r_s}$, $r_s = \frac{1}{2} \sum_j k_j$ being the reduction power:

$$\otimes_{j=1}^s \bar{\mathcal{M}}_{g_j, n_j + k_j}^{disc} (-1)^{r_s}. \quad (4.3)$$

Integration over $\bar{\mathcal{M}}_{g_j, n_j + k_j}^{disc} (-1)^{r_s}$ expands into product of integrals over connected closed components which are given by known continuous Kontsevich’s indices. So we conjecture the answer for intersection indices on the discretized moduli space:

$$\frac{2^d}{d!} \int_{\mathcal{M}_{g,n}^{disc}} \Omega^d = \sum_{\sum d_i = d} \prod_{i=1}^n \frac{p_i^{2d_i}}{d_i!} \langle \tau_{d_1} \dots \tau_{d_n} \rangle_g$$

$$+ \sum_{\text{reductions}} (-1)^{r_s} \frac{(d-r_s)!}{d!} \prod_{j=1}^s \left(\sum_{d_a=d_j} \prod_{a=1}^{n_j} \frac{p_a^{2d_a}}{d_a!} < \tau_{d_1} \dots \tau_{d_{n_j}} \tau_0^{(1)} \dots \tau_0^{(k_j)} >_{g_j} \right), \quad (4.4)$$

where $< \dots >_{g_j}$ are corresponding Kontsevich's indices, $d_j = 3g_j - 3 + n_j + k_j$ and k_j insertions of τ_0 correspond to additional punctures.

Next step is to do the Laplace transform over variables p_i . Now we should sum over $p_i = \epsilon, 2\epsilon, 3\epsilon, \dots$ simultaneously taking into account that $\sum_{i=1}^n p_i \in 2\epsilon\mathbf{Z}_+$ (every edge n_a is counted twice). A procedure is the following. We do the transform over variables λ_j taking a sum over all $p_j \in \epsilon\mathbf{Z}_+$ weighted with $(i)^j$ and release the real part of the obtained expression consequently substituting $e^{\epsilon\lambda_j} \rightarrow i e^{\epsilon\lambda_j}$. On the right hand side we have (explicitly reconstructing the ϵ -dependence):

$$\text{Re} \prod_{j=1}^n \left(\sum_{p_j=1}^{\infty} e^{-\epsilon\lambda_j p_j} (i)^{p_j} \epsilon^{2d_j} p_j^{2d_j} \right) = \prod_{j=1}^n \left(\frac{\partial}{\partial \lambda_j} \right)^{2d_j} \text{Re} \frac{(-1)^n}{\prod_{j=1}^n (1 + i e^{\epsilon\lambda_j})}. \quad (4.5)$$

Taking as an example the case of $\mathcal{M}_{0,3}$ we immediately get the expression (3.5).

On the right hand side we get:

$$\begin{aligned} & \sum_{\{p_i\} \in \mathcal{M}_{g,n}^{disc}} e^{-\epsilon\lambda_j p_j} \epsilon^n e^{\Omega} dp_1 \wedge \dots \wedge dp_n \\ &= \int_{\mathcal{M}_{g,n}^{disc}} e^{-\epsilon\lambda_j p_j} 2^{2g-2} dn_1 \wedge \dots \wedge dn_{6g-6+3n} \epsilon^{6g-6+3n}. \end{aligned} \quad (4.6)$$

The last term possesses an explicit representation as a sum over all possible graphs with fixed genus g and number of cycles n and *arbitrary* valencies of vertices:

$$\begin{aligned} & \sum_{\Gamma} \frac{1}{\# \text{Aut}(\Gamma)} 2^{\# \text{edges} - \# \text{vert.} + n} \epsilon^{6g-6+3n} \prod_s \sum_{n_s=1}^{\infty} e^{-\epsilon n_s (\lambda_s^{(1)} + \lambda_s^{(2)})} \\ &= 2^n \sum_{\Gamma} \frac{1}{\# \text{Aut}(\Gamma)} 2^{-\# \text{vert.}} \epsilon^{6g-6+3n} \prod_{\{ij\}} \frac{2}{e^{\epsilon(\lambda_i + \lambda_j)} - 1} \\ &\equiv w_g(\lambda_1, \dots, \lambda_n). \end{aligned} \quad (4.7)$$

This last expression is in fact the free energy term for the Kontsevich–Penner matrix model in fixed g and n of the form

$$e^{F_N(\Lambda)} = \frac{\int DX \exp N\alpha \text{tr} \left\{ -\frac{1}{4} \Lambda X \Lambda X - \frac{1}{2} [\log(1-X) + X] \right\}}{\int DX \exp N\alpha \text{tr} \left\{ -\frac{1}{4} \Lambda X \Lambda X + \frac{1}{4} X^2 \right\}}, \quad (4.8)$$

where $\alpha = 1/\epsilon^3$ and $\Lambda = \text{diag}(e^{\epsilon\lambda_1}, e^{\epsilon\lambda_2}, \dots, e^{\epsilon\lambda_N})$.

$F_N(\Lambda)$ has an expansion which looks just like (1.3):

$$F_N(\Lambda) = \sum_{\substack{g=0 \\ n=1}}^{\infty} (N\alpha)^{2-2g} \alpha^{-n} N^{-n} \text{tr} w_g(\lambda_1, \dots, \lambda_n). \quad (4.9)$$

In the continuum limit $\epsilon \rightarrow 0$ we immediately get the expression (2.2) i.e. the Kontsevich matrix model.

Thus we have demonstrated the relation between geometric characteristics of the discretized moduli space and related matrix models. We show how the Kontsevich indices can be naturally embedded into standard one-matrix model via the Penner–Kontsevich model.

In conclusion we should note some problems and perspectives of the proposed model. First, we did not yet prove rigorously the relation (4.4), and this proof is necessary for the completeness of the theory. Also it is interesting to make the sum over reductions more explicitly (in the combinatorial sense). Second interesting problem to solve is to find the description of just the Penner model in the case of the discretized moduli space. The formula (3.2) should be modified since in this case the volume of the stabilizer (symmetry group) taken at the boundary (infinite) point might be no infinite but rather proportional to some positive power of ϵ , the discretization parameter. Also it is interesting to develop this approach to the case of GKM, where most subtle geometric invariants are considered.

I am grateful to G.Falqui, V.Fock, A.Gerasimov, Yu.Makeenko and A.Mironov for numerous valuable discussions.

References

- [1] E.Br ezin and V.Kazakov, *Phys. Lett.* **236B** (1990) 144;
M.Douglas and S.Shenker, *Nucl. Phys.* **B335** (1990) 635;
D.J.Gross and A.Migdal, *Phys. Rev. Lett.* **64** (1990) 127.
- [2] A.Mironov and A.Morozov, *Phys. Lett.* **B252** (1990) 47;
J.Ambj rn, J.Jurkiewicz, and Yu.Makeenko, *Phys.Lett.*, **251B** (1990) 517;
A.Gerasimov, A.Marshakov, A.Mironov, A.Morozov, and A.Orlov, *Nucl.Phys.*, **B357** (1991) 565;
H.Itouyama and Y.Matsuo, *Phys.Lett.*, **B255** (1991) 202;
Yu.Makeenko, A.Marshakov, A.Mironov and A.Morozov, *Nucl. Phys.* **B356** (1991) 574;
H.S.La, *Commun. Math. Phys.* **140** (1991) 569;
M.Fukuma, H.Kawai and R.Nakayama, *Infinite dimensional Grassmannian Structure of two-dimensional Quantum Gravity*, University of Tokyo preprint UT-572 (November, 1990).
- [3] E.Witten *Nucl. Phys.* **B340** (1990) 281.
- [4] E.Witten *Surveys in Diff. Geom.* **1** (1991) 243.
- [5] M.L.Kontsevich, *Funk.Anal.&Prilozh.*, **25** (1991) 50 (in Russian); *Intersection Theory on the Moduli Space of Curves and the Matrix Airy Function*, Max-Planck-Institute preprints MPI/91-47 and MPI/91-77.

- [6] M.Fukuma, H.Kawai and R.Nakayama, *Int. J. Mod. Phys.* **A6** (1991) 1385;
R.Dijkgraaf, E.Verlinde and H.Verlinde, *Nucl. Phys.* **B348** (1991) 435.
- [7] E.Witten, *On the Kontsevich Model and other Models of Two-Dimensional Gravity*,
Princeton preprint IASSNS-HEP-91/24 (June, 1991).
- [8] D.J.Gross and M.J.Newman, *Phys.Lett.*, **266B** (1991) 291; *Unitary and hermitian
matrices in an external field II: the Kontsevich model and continuum Virasoro con-
straints*, preprint *PUP-T-1282* (December, 1991).
- [9] A.Marshakov, A.Mironov, and A.Morozov, *On equivalence of topological and quantum
2d gravity*, preprint *FIAN/TD/04-91* (August, 1991).
- [10] R.Dijkgraaf, *Intersection theory, integrable hierarchies and topological field theory*,
preprint *IASSNS-HEP-91/91* (December, 1991).
- [11] E.Witten, *The N matrix model and gauga WZW models*, preprint *IASSNS-HEP-
91/26* (June, 1991); *Algebraic geometry associated with matrix models of two dimen-
sional gravity*, preprint *IASSNS-HEP-91/74* (October, 1991);
R.Dijkgraaf and E.Witten, *Nucl. Phys.* **B342** (1990) 486;
K.Li, *Nucl. Phys.* **B354** (1991) 711, 725
R.Dijkgraaf, E.Verlinde and H.Verlinde, *Nucl. Phys.* **B352** (1991) 59;
S.Kharchev, A.Marshakov, A.Mironov, A.Morozov, and A.Zabrodin, *Unification of
all string models with $c < 1$; Towards unified theory of 2d gravity*, *FIAN/ITEP
preprints* (October, 1991).
- [12] R.C.Brower and M.Nauenberg, *Nucl.Phys.*, **B180[FS2]** (1981) 221;
E.Br zin and D.Gross, *Phys.Lett.*, **97B** (1980) 120;
R.C.Brower, P.Rossi, and C.-I Tan, *Phys.Rev.*, **D23** (1981) 942.
- [13] Yu.Makeenko and G.Semenoff, *Mod.Phys.Lett.*, **A6** (1991) 3455;
Yu.Makeenko, *Mod.Phys.Lett. (Brief Reviews)*, **A6** (1991) 1901.
- [14] C.Itzykson and J.-B.Zuber, *Combinatorics of the modular group II. The Kontsevich
integrals*, preprint *SPhT/92-001* (January, 1992).
- [15] L.Chekhov and Yu.Makeenko, *The multicritical Kontsevich-Penner model*, preprint
NBI-HE-92-03 (January, 1992).
- [16] J.Amb rn, L.Chekhov and Yu.Makeenko, *Higher Genus Correlators and W-infinity
from the Hermitian One-Matrix Model*, preprint *NBI-HE-92-22* (February, 1992).
- [17] L.Chekhov and Yu.Makeenko, *Hint on the External Field Problem for Matrix Models*,
preprint *NBI-HE-92-06* (January, 1992).
- [18] S.Kharchev, A.Marshakov, A.Mironov, and A.Morozov, *Generalized Kontsevich
Model versus Toda hierarchy*, *Lebedev Institute preprint FIAN/TD-03/92* (febru-
ary, 1992).

- [19] P.Deligne and M.Mumford *The irreducibility of the space of Curves of given Genus* *Publ. I.H.E.S.* **45** (1969) 75.
D.Mumford, *Towards an Enumerative Geometry of the Moduli Space of Curves*, in *Arithmetic and Geometry*, vol II, Birkhäuser, 1983.
- [20] J.Harer and D.Zagier, *Invent. Math.*, **85** (1986) 457;
J.Harer, *The Cohomology of the Moduli Space of Curves* (Gordon, 1985).
- [21] R.C.Penner, *The Decorated Teichmüller Space of Punctured Surfaces*, *Comm. Math. Phys.* **113** (1987), 299–339; *Perturbative Series and the Moduli Space of Riemann Surfaces*, *J. Diff. Geom.* **27** (1988), 35–53.
- [22] K.Strebel, “Quadratic Differentials”, Springer–Verlag, 1984.
- [23] I.K.Kostov and M.L.Metha, *Phys.Lett.*, **189B** (1987) 118.
- [24] J.Distler and C.Vafa, *Mod.Phys.Lett.*, **A6** (1991) 259.